



Additional solutions of a weighted heat equation

Da-qian Lu

East China Normal University, Department of Mathematics, Shanghai 200241, PR China

ARTICLE INFO

Article history:

Received 5 May 2010

Accepted 18 May 2011

Keywords:

Weighted heat equation

Legendre polynomial

Additional solutions

ABSTRACT

We obtain a simple algorithm for computing additional solutions of a weighted heat equation.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

We recall Legendre's differential equation [1]

$$\frac{d}{dx} \left[(1-x^2) \frac{dv(x)}{dx} \right] + n(n+1)v(x) = 0 \quad (1.1)$$

which has the Legendre polynomial given by Rodrigues' formula:

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n [(x^2-1)^n]}{dx^n} \quad (1.2)$$

as one of its solutions.

The series representation is

$$P_n(x) = \sum_{k=0}^{[n/2]} \frac{(-1)^k (1/2)_{n-k} (2x)^{n-2k}}{k!(n-2k)!} \quad (1.3)$$

with the generating function

$$\frac{1}{\sqrt{1-2xt+t^2}} = \sum_{n=0}^{\infty} P_n(x)t^n. \quad (1.4)$$

Expanding the Taylor series in Eq. (1.4) for the first two terms gives

$$P_0(x) = 1, \quad P_1(x) = x$$

for the first two Legendre polynomials. To obtain further terms, Eq. (1.4) is differentiated with respect to t on both sides and rearranged to give

$$\frac{x-t}{\sqrt{1-2xt+t^2}} = (1-2xt+t^2) \sum_{n=0}^{\infty} nP_n(x)t^{n-1}.$$

E-mail address: daqianlu@163.com.

Replacing the quotient of the square root with its definition in (1.4), and equating the coefficients of powers of t in the resulting expansion gives Bonnet's recursion formula:

$$(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x). \quad (1.5)$$

This relation, along with the first two polynomials P_0 and P_1 , allows the Legendre polynomials to be generated recursively. The first few Legendre polynomials are

$$\begin{aligned} P_2(x) &= \frac{3}{2}x^2 - \frac{1}{2}, & P_3(x) &= \frac{5}{2}x^3 - \frac{3}{2}x, \\ P_4(x) &= \frac{35}{8}x^4 - \frac{15}{4}x^2 + \frac{3}{8}, & P_5(x) &= \frac{63}{8}x^5 - \frac{35}{4}x^3 + \frac{15}{8}x. \end{aligned} \quad (1.6)$$

In [2], the author considered the weighted heat equation

$$\frac{d}{dx} \left[w(x) \frac{dv(x)}{dx} \right] = -\lambda^2 v(x) \quad (1.7)$$

in the special case of

$$\frac{d}{dx} \left[(1-x^2) \frac{dv(x)}{dx} \right] = -\lambda^2 v(x). \quad (1.8)$$

Then Eq. (1.8) is Legendre's differential equation when $\lambda^2 = n(n+1)$.

Suppose that $v_{1,n} = P_n(x)$; then from the variation of parameters method, the additional solutions are given by the formula

$$v_{2,n} = P_n(x)q_n(x) \quad (1.9)$$

where

$$q_n(x) = \int \frac{dx}{(1-x^2)P_n^2(x)}. \quad (1.10)$$

From (1.10), we have

$$q_0(x) = \int \frac{dx}{1-x^2} = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right|.$$

Then, by (1.9) we get

$$v_{2,0} = P_0(x)q_0(x) = \ln \sqrt{\left| \frac{1+x}{1-x} \right|}.$$

Using the same method, another three solutions are calculated:

$$\begin{aligned} v_{2,1} &= x \ln \sqrt{\left| \frac{1+x}{1-x} \right|} - 1, \\ v_{2,2} &= \frac{1}{2}(3x^2 - 1) \ln \sqrt{\left| \frac{1+x}{1-x} \right|} - \frac{3}{2}x \end{aligned}$$

and

$$v_{2,3} = \frac{1}{2}(5x^3 - 3x) \ln \sqrt{\left| \frac{1+x}{1-x} \right|} - \frac{5}{2}x^2 + \frac{2}{3}.$$

Continued calculations carried out in a manner similar to those above would generate an infinite number of solutions to (1.8). But the integration in (1.10) becomes more and more complex as the index n increases.

In this work, we get a simple algorithm which allows us to do without the prohibitive integrations necessary to compute additional solutions.

2. The main result

Theorem 2.1. *The solutions of (1.8) satisfy the following relation:*

$$v_{2,n+1}(x) = \frac{2n+1}{n+1}x v_{2,n}(x) - \frac{n}{n+1} v_{2,n-1}(x). \quad (2.1)$$

Proof. We assume that the relation (2.1) is satisfied for $k = n$, that is

$$v_{2,n}(x) = \frac{2n-1}{n}x v_{2,n-1}(x) - \frac{n-1}{n} v_{2,n-2}(x). \quad (2.2)$$

Then, using (1.9), we have

$$nP_n(x)q_n(x) = (2n-1)xP_{n-1}(x)q_{n-1}(x) - (n-1)P_{n-2}(x)q_{n-2}(x). \quad (2.3)$$

If in (1.5) we decrease n by 1 we have

$$nP_n(x) = (2n-1)xP_{n-1}(x) - (n-1)P_{n-2}(x). \quad (2.4)$$

Substituting the recurrence relation (2.4) into Eq. (2.3) we find

$$(2n-1)xP_{n-1}(x)q_n(x) - (n-1)P_{n-2}(x)q_n(x) = (2n-1)xP_{n-1}(x)q_{n-1}(x) - (n-1)P_{n-2}(x)q_{n-2}(x). \quad (2.5)$$

If in (2.5) we increase n by 1 we get

$$(2n+1)xP_n(x)q_{n+1}(x) - nP_{n-1}(x)q_{n+1}(x) = (2n+1)xP_n(x)q_n(x) - nP_{n-1}(x)q_{n-1}(x). \quad (2.6)$$

By using (1.5) and (1.9) we can get

$$\begin{aligned} v_{2,n+1}(x) &= P_{n+1}(x)q_{n+1}(x) \\ &= \frac{2n+1}{n+1}xP_n(x)q_{n+1}(x) - \frac{n}{n+1}P_{n-1}(x)q_{n+1}(x). \end{aligned} \quad (2.7)$$

Substituting (2.6) into (2.7) we have

$$v_{2,n+1}(x) = \frac{2n+1}{n+1}xP_n(x)q_n(x) - \frac{n}{n+1}P_{n-1}(x)q_{n-1}(x). \quad (2.8)$$

Using (1.9) again we get the relation (2.1). \square

Remark 2.1. It is interesting to note from the above theorem that each solution is dependent on the two preceding solutions. So we can compute the additional solutions of (1.8) easily; for example,

$$\begin{aligned} v_{2,4} &= \frac{1}{8}(35x^4 - 30x^2 + 3) \ln \sqrt{\left| \frac{1+x}{1-x} \right|} - \frac{35}{8}x^3 + \frac{55}{24}x, \\ v_{2,5} &= \frac{1}{8}(63x^5 - 70x^3 + 15x) \ln \sqrt{\left| \frac{1+x}{1-x} \right|} - \frac{63}{8}x^4 + \frac{49}{8}x^2 - \frac{8}{15}, \end{aligned}$$

etc.

Acknowledgments

The author would like to thank the referees and editors for their many valuable comments and suggestions. This work was supported by the Shanghai Natural Science Foundation, China under Grant 10ZR1409100.

References

- [1] E.D. Rainville, Special Functions, The Macmillan Company, New York, 1960.
- [2] G.A. Bear, Legendre functions as solutions to the inhomogeneous heat equation, A Thesis in Mathematics, 1995.